# Sexual Function and Mental Health Medications

## Common Questions, Real Effects, and Practical Options

### Why do mental health medications affect sexual function?

Sexual response depends on a delicate balance between the nervous system, hormones, blood flow, and psychological factors. Mental health medications act on brain chemicals (neurotransmitters) that regulate mood, anxiety, and behaviour, but these same systems are also involved in sexual desire, arousal, and orgasm.
Because of this overlap, sexual side effects are common and do not mean that treatment is failing or that something is "wrong" with the patient.

### Which phases of sexual function can be affected?

Mental health medications can influence different phases of sexual activity, including:

- **Sexual desire** (reduced interest in sex)
- **Arousal** (difficulty becoming or staying aroused)
- **Orgasm** (delayed or absent orgasm)
- **Overall sexual satisfaction**

The type and severity of effects vary widely between individuals and medications.

### How do antidepressants affect sexual health?

Antidepressants, especially **SSRIs** and **SNRIs**, are commonly associated with sexual dysfunction.
They may:

- Reduce libido
- Delay ejaculation or orgasm
- Make orgasm more difficult or absent

For some people, these effects are mild. For others, they can be distressing and may affect adherence to treatment.

### Can sexual side effects ever be helpful?

In certain situations, yes. For example:

- Some men with **premature ejaculation** may benefit from delayed ejaculation caused by SSRIs.
- SSRIs are sometimes used to manage **compulsive sexual behaviours**, because reducing libido can lower distressing urges.

This highlights that sexual effects are not always purely negative, but context matters.

### Are there antidepressants with fewer sexual side effects?

Yes. Some antidepressants are associated with **lower rates of sexual dysfunction**, including:

- **Bupropion**

- **Mirtazapine**
- **Trazodone**

In selected cases, switching medications can improve sexual function without compromising mental health treatment, as shown in clinical studies (for example with mirtazapine).

### What about anxiety medications?

Anxiety disorders often require antidepressant treatment, which can also affect sexual function.

- **SSRIs** used for anxiety commonly reduce sexual desire and may cause ejaculation difficulties.
- **Benzodiazepines**, sometimes prescribed short term, tend to have fewer direct sexual side effects, although they carry other risks and are not a long-term solution.

It is often difficult to separate the effects of anxiety itself from those of medication.

### How do antipsychotic medications affect sexuality?

In psychotic disorders, **antipsychotic medications** are a major cause of sexual dysfunction.
Common effects include:

- Reduced libido
- Difficulties with arousal
- Problems with orgasm

Some antipsychotics (including older drugs and **risperidone**) can increase **prolactin levels**, which interferes with sexual hormones and function.
Newer medications, such as **aripiprazole**, may reduce these effects in some patients and can sometimes improve sexual function.

### Can these medications affect sexual health indirectly?

Yes. Some medications influence metabolism, weight, and energy levels. These changes can affect sexual function:

- **Directly**, through physical health changes
- **Indirectly**, by reducing vitality or encouraging a sedentary lifestyle

Lifestyle support and education can therefore be an important part of care.

### What is the role of healthcare professionals?

Sexual health is a key part of quality of life and should be addressed routinely. Clinicians should help clarify:

- Did sexual difficulties start **before** or **after** treatment?
- Are they linked to the mental health condition, the medication, or both?
- How do symptoms change over time?

Management may include medication adjustments, psychosexual education, lifestyle support, and reframing expectations toward a **"good enough sex"** model that fits the person's needs and context.

> ***Key Message***
>
> *Mental health medications can affect sexual function, but this does not mean sexual wellbeing must be sacrificed. Asking the right questions and reviewing treatment options can help balance mental health stability with a satisfying sexual life.*

Credits: Giedre Jonusiene (Lithuania)
Wojciech Merk (Poland)